



University of Mazandaran

JGR

Journal of Genetic Resources

Vol. 1, No. 1 (2015): 7-18

<http://sc.journals.umz.ac.ir>



Iranian Biology Society

Influence of Air SO₂ Pollution on Antioxidant Systems of Alfalfa Inoculated with Rhizobium

Shima Hosseinkhani Hezave¹, Mehri Askari^{2*}, Fariba Amini², Morteza Zahedi³

¹ MSc. of Plant Physiology, Biology Department, Faculty of Science, Arak University

² Biology Department, Faculty of Science, Arak University, Arak 38156-8-8349, Iran

³ Amirkabir Technical and Vocational College, Arak

* Corresponding author: m-askary@araku.ac.ir

Received: 12/08/2013 Accepted: 10/01/2014

Abstract

Air pollution is the most important environmental problem of last century that threatens the health of living organisms, especially plants. SO₂ is one of the main air pollutants that can cause to imbalance in growth and physiological function of plant in high concentrations. Symbiosis of *Rhizobium* bacteria with alfalfa can cause increasing plant growth and resistance to abiotic stresses. In order to study the effects of rhizobia inoculation on alfalfa antioxidant activity and capacity under air SO₂ pollution, 35 days plants (noninoculated and inoculated with native or standard *Rhizobium meliloti*) exposed to the different concentrations of SO₂ (0 as a control, 0.5, 1, 1.5 and 2 ppm) for 6 consecutive days (2 hours per day). Results showed inoculation had no significant effect on antioxidant activity and capacity of alfalfa plant. However different concentrations of SO₂ pollution had a significant effect on alfalfa antioxidant system. Increasing SO₂ stress increased antioxidant activities (I%) and decreased antioxidant capacities (IC₅₀) of alfalfa leaves significantly in comparison to the control plants (under 0 ppm) as well as increased superoxide dismutase, catalase and guaiacol peroxidase activity. Inoculation of alfalfa plant with *Rhizobium meliloti* reduced the negative effects of high concentrations of SO₂ on antioxidant activity and capacity significantly. In fact, the rate of this change of antioxidant system under SO₂ pollution in inoculated plants was lower than in the noninoculated plants. Therefore inoculation with *Rhizobium* strains could alleviate the effect of SO₂ pollution on antioxidant system.

Keywords: Alfalfa, Antioxidant activity and capacity, *Rhizobium*, SO₂ pollution

تأثیر آلودگی SO₂ هوا بر سیستم آنتی‌اکسیدانی یونجه تلقیح یافته با ریزوبیوم

شیمای حسین‌خانی هزاوه¹، مه‌ری عسکری²، فریبا امینی²، مرتضی زاهدی³

¹ دانشجوی کارشناسی ارشد فیزیولوژی گیاهی، دانشکده علوم، دانشگاه اراک

² دانشگاه اراک، دانشکده علوم، گروه زیست‌شناسی، کد پستی 38156-8-8349

³ دانشگاه فنی و حرفه‌ای امیرکبیر اراک

چکیده

آلودگی هوا از مهمترین مشکلات زیست محیطی قرن اخیر است که سلامت موجودات زنده به‌ویژه گیاهان را تهدید می‌نماید. SO₂ یکی از مهمترین آلاینده‌های هوا است که در غلظت‌های بالا می‌تواند باعث اختلال در رشد و عملکرد فیزیولوژیکی گیاهان شود. همزیستی باکتری ریزوبیوم با یونجه می‌تواند باعث افزایش رشد و مقاومت گیاه نسبت به تنش‌های غیر زیستی شود. به‌منظور مطالعه اثرات تلقیح ریزوبیوم بر ظرفیت و فعالیت آنتی‌اکسیدانی یونجه تحت آلودگی SO₂، گیاهان 35 روزه (تلقیح نیافته و تلقیح یافته با *Rhizobium meliloti* استاندارد یا بومی) به‌مدت 6 روز متوالی (روزی 2 ساعت) در معرض غلظت‌های مختلف SO₂ (0 به عنوان شاهد، 0.5، 1، 1.5 و 2 ppm) قرار گرفتند. نتایج نشان داد تلقیح اثر معنی‌داری روی ظرفیت و فعالیت آنتی‌اکسیدانی گیاه یونجه نداشت. گرچه غلظت‌های مختلف آلودگی SO₂ اثر معنی‌داری روی سیستم آنتی‌اکسیدانی یونجه داشت. افزایش تنش SO₂ به‌طور معنی‌داری باعث افزایش فعالیت آنتی‌اکسیدانی (I%) و کاهش ظرفیت آنتی‌اکسیدانی (IC₅₀) برگ‌های یونجه در مقایسه با گیاهان شاهد (تحت 0ppm) شد همچنین فعالیت سوپراکسید دیسموتاز، کاتالاز و گایاکول پراکسیداز افزایش یافت. تلقیح گیاه یونجه با باکتری *Rhizobium meliloti* اثرات منفی غلظت‌های بالای SO₂ روی ظرفیت و فعالیت آنتی‌اکسیدانی را به‌طور معنی‌داری کاهش داد. در واقع نرخ تغییر سیستم آنتی‌اکسیدانی تحت آلودگی SO₂ در گیاهان تلقیح یافته کمتر از گیاهان تلقیح نیافته بود. بنابراین تلقیح با سوبه‌های ریزوبیوم می‌تواند اثرات آلودگی SO₂ روی سیستم آنتی‌اکسیدانی را کم کند.

واژه‌های کلیدی: آلودگی SO₂؛ ریزوبیوم؛ ظرفیت و فعالیت آنتی‌اکسیدانی؛ یونجه

Introduction

Air pollution has become an extremely serious problem for the modern industrialized world. Air pollution may be defined as any atmospheric condition which certain substances are present in concentrations that may produce undesirable effects on human and ecosystem. These substances include gases (SO₂, nitrogen oxides, carbon monoxides, and hydrocarbons), particulate matters (smoke, dust, fumes, and aerosols) and radioactive materials (Gostin, 2009; Rai *et al.*, 2011). Air pollution was earlier considered as a local problem around large point sources. But due to use of tall stacks and long range transport of pollutants, it has become a regional problem. Uncontrolled use of fossil fuels in industries and transport sectors has led to the increase in concentrations of gaseous pollutants such as SO₂, NO_x (Rai *et al.*, 2011).

Sulfur dioxide (SO₂) is one of the most common and harmful air pollutants (Li & Yi, 2012) that its concentration is increasing in many metropolitan and industrial areas. It is a major atmospheric contaminant resulting primarily from the combustion of sulphur-rich fossil fuels such as coal and oil or naturally from forest fire and volcanic eruptions. In the atmosphere, when SO₂ combines with water, it forms sulphurous acid which is the main component of acid rain (Sha *et al.*, 2010). SO₂, normally 0.05–0.5 ppm in the urban areas and up to 2 ppm or more around sources of air pollution (Wali *et al.*, 2007). SO₂ and acid rain be harmful to plants, turning leaves yellow and dry, bleaching and even killing foliage depending on the dosages (Lang *et al.*, 2007). The 40% of SO₂ global emissions originating from Asia and it is growing (Smith *et al.*, 2011). The phytotoxicity of SO₂ strongly depends on its concentration and exposure duration and is also influenced by the sulfur status of plants (Li & Yi, 2012).

In very low concentrations, SO₂ can cause positive effects on physiological and growth of plants, especially in sulfur-deficient soils (Swanepoel *et al.*, 2007), since sulfur is a structural component of amino acids, proteins, vitamins and chlorophyll. Sulfur enhances the development of nodules and nitrogen fixation by legumes and also affects carbohydrate metabolism (Li & Yi, 2012). However, exposure to high doses of SO₂ causes toxicity and reduction of growth due to sulphite and sulphate accumulation within

the plant (Swanepoel *et al.*, 2007). SO₂ can easily penetrate into chloroplasts and affect plant growth and development (Sha *et al.*, 2010). Even when stomata are closed, SO₂ can react with water to produce bisulfite and enter the leaf through the cuticle (Sha *et al.*, 2010). In the chloroplasts, SO₂ is mainly converted into sulfite, which causes a reduction of net CO₂ assimilation, inhibits photosynthetic enzymes, and decreases the photosynthetic electron transport rate (Sha *et al.*, 2010). SO₂-toxicity is mainly attributed to produce high reactive intermediates such as the sulphur trioxide radical (HSO₃[·]), the superoxide radical and the hydroxyl radical, which are generated during the radical-initiated oxidation of SO₂. Illuminated chloroplasts can initiate oxidation of SO₂ and, hence, may be a primary site of radical production during SO₂ treatment. To counteract the toxicity of active oxygen species, a high efficient antioxidative defence system, composed of both non-enzymatic (e.g. glutathione, proline, α-tocopherols, carotenoids and flavonoids) and enzymatic (e.g. superoxide dismutase, catalase, glutathione peroxidase and glutathione reductase) constituents is present in all plant cells (Gill & Tuteja, 2010; Li & Yi, 2012).

Rhizobium is one of the most prominent Plant growth promoting rhizobacteria (PGPR) members that activate plant root nodulation in leguminous plants (Gentili & Jumpponen, 2006). PGPR can promote plant growth by several mechanisms, directly and/or indirectly, include (i) the ability to produce or change the concentration of plant growth regulators like indoleacetic acid, gibberellic acid, cytokinins and ethylene, (ii) N₂ fixation, (iii) production of siderophores, antibiotics and cyanide (iv) solubilization of mineral phosphates and other nutrients (Aeini *et al.*, 2012; Martínez-Viveros *et al.*, 2010; Saharan & Nehra, 2011). PGPR can prevent the deleterious effects of environmental stressors (Han & Lee, 2005; Tank & Saraf, 2010). The use of PGPR offers an attractive way to replace chemical fertilizer, pesticides and supplements. Some PGPR have been produced commercially as inoculants for agriculture to improve plant growth through supply of plant nutrients and may help to sustain environmental health and soil productivity (Estefan *et al.*, 2013; Tank & Saraf, 2010).

Alfalfa (*Medicago sativa* L.) is a very used forage legume with over 32 million hectares in the world (Benabderrahim *et al.*, 2009) and it's the most important forage crop for the arid and semi-arid

areas (Salehi *et al.*, 2008). The objective of this study was to determine the effects of rhizobium inoculation on antioxidant activity and total antioxidant of alfalfa under different concentrations of SO₂ (0, 0.5, 1, 1.5 and 2 ppm).

Materials and Methods

Bacterial culture and inoculant preparation

Alfalfa plants were collected from Arak farm lands (a city of Iran that high levels of SO₂ has been reported (Moini *et al.*, 2011) and native strain of *Rhizobium meliloti* was extracted from roots of this plants. For this purpose, alfalfa roots were sterilized with 70% ethanol and were washed with sterile distilled water (Swift & Bignell, 2001). Then the pink nodules (containing active bacteria) isolated from roots, crushed in distilled water and cultured in solid medium of YMA (Yeast Manitol Agar) (Molla *et al.*, 2001). These cultures were transferred to incubator at 25°C. After incubation, gram reaction and morphology of bacteria were studied under the microscope. Formation of convex prominent semi-transparent slimy and mucilage colonies and gram-negative reaction were considered a sign of successful isolation of rhizobium (Swift & Bignell, 2001).

Standard strain of rhizobium (*Rhizobium meliloti* PTCC 1684) were obtained from the Persian type culture collection (PTCC, Iran). For activation these bacteria, 1 ml of liquid medium of YMA under sterile conditions added to powdered of bacteria. For proliferation of bacteria, one inoculation loop of these bacteria dissolved in 100 ml of liquid YMA and incubated on an orbital shaker at 200 rpm for 24 h. Optimum amount of rhizobium to stimulate alfalfa growth was reported 10⁵ cells/mL (Caetano-Anolles *et al.*, 1988), for this purpose two strains of *Rhizobium meliloti* (native and standard) were cultured separately in liquid medium of YMA (Molla *et al.*, 2001) and incubated on an orbital shaker at 200 rpm for 24 h at 25°C (Sadovinkova *et al.*, 2003). Then these cultures were centrifuged at 1000g for 10 min and were resuspended with phosphate buffer. The optical density (OD₆₂₀) of this solution was 0.1 it means 10⁸ cells/mL (Bai *et al.*, 2003). For preparation optimum amount of inoculum (10⁵ cells/mL), this solution was diluted by phosphate buffer.

Seed preparation and its inoculation

The seeds of alfalfa (*Medicago sativa* cv. Hamedani) were prepared from Arak Agriculture Research Center. They were sterilized by 70% ethanol for 2 min and 1% sodium hypochlorite for 5 min, then they washed with distilled water 5 times (Wang & Oyaizu, 2009). After washing, the seeds were divided into three groups. First group of seeds inoculated with native *R. meliloti* inoculum, second group of seeds inoculated with standard *R. meliloti* inoculum and third group soaked in sterile phosphate buffer. All of the groups were placed under vacuum and ambient temperature for 2 h (Bashan *et al.*, 1989).

Hydroponic cultivation of seed

Inoculated and non-inoculated seeds were placed to plates system containing nutrient solution (without nitrogen) in the dark for 24 h. The germinated seeds were transferred to sterile microtubes in plastic container containing 2 L of nutrient solution without nitrogen (Millner & Kitt, 1992). Containers were oxygenated by the air compressor. Each container was considered as an experimental group. These containers were maintained under 12 h photoperiod, at 25°C during day and 20°C during night (Bashan *et al.*, 1989).

SO₂ treatments for plant

SO₂ gas prepared from Shazand Petrochemical Co. injected in different concentrations 0 (as control), 0.5, 1, 1.5 and 2 ppm into 35 days plants. Gas injection was performed by syringe for 5 days and 2 h daily to closed plastic containers (Agrawal *et al.*, 1985).

Enzyme assays

Extraction: Leaf fresh materials (0.1 g) was powdered by liquid nitrogen and homogenized in 1 ml of 50 mM phosphate buffer (pH 7) containing 1 mM ethylene diamine tetra acetic acid (EDTA) by a homogenizer into microtubes. Insoluble materials removed by Beckman refrigerated centrifuge at 13000 g for 20 min at 4°C, and the supernatant used as the source of enzyme extraction.

Assays: All enzyme assays were performed at 4°C and activities of the enzymes were determined with a spectrophotometer (PG T80 UV/VIS, Oasis Scientific Inc.).

Superoxide dismutase (SOD) assay Superoxide dismutase activity was measured by monitoring

the inhibition of photochemical reduction of nitro blue tetrazolium according to the method described by Giannopolitis & Ries (1977), using a reaction mixture (3 mL) containing 50 mM phosphate buffer (pH = 7.8), 13 mM methionine, 75 μ M nitro blue tetrazolium, 20 μ M riboflavin, 0.1 mM EDTA and 100 μ l of the enzyme extract in absence of light. The reaction mixtures were illuminated for 15 min under fluorescent light. One unit of superoxide dismutase activity is defined as the amount of enzyme required to cause 50% inhibition of nitroblue tetrazolium reduction, which was monitored at 560 nm.

Catalase (CAT) assay

Catalase activity was assayed by measuring the initial amount of H₂O₂ disappearance using the method of Cakmak & Marschner (1992) in a reaction mixture containing 2 ml of 25 mM phosphate buffer (pH 7.0, containing 10 mM H₂O₂) and 100 μ l of the enzyme extract. Decomposition of 1 μ mol H₂O₂/min is equal to one unit of catalase activity.

Guaiacol peroxidase (GPOX) assay

Guaiacol peroxidase activity was measured by Polle *et al.* (1994). The reaction mixture contained 100 mM phosphate buffer (pH 7.0), 20 mM guaiacol, 10 mM H₂O₂ and 50 μ l of the enzyme extract. Activity was determined by increasing absorbance at 470 nm due to guaiacol oxidation.

Measurement of DPPH-radical scavenging activity

For determination of DPPH (1, 1-diphenyl-2-picrylhydrazyl) radical scavenging activity used of Abe *et al.* (1998) method. Leaf fresh materials (100 mg) was powdered by liquid nitrogen, homogenized in 1 ml of 90% ethanol and then maintained at 4°C for 24 h. Insoluble materials removed by centrifuge at 3500 g for 5 min. 20 μ l of extracting solution was mixed with 800 μ l of DPPH (0.5 mM in ethanol). The absorbance of the resulting solution was measured at 517 nm after 30 min in darkness. The antiradical capacity (three replicates per treatment) was expressed as IC₅₀ (mg ml⁻¹), the antiradical dose required to cause a 50% inhibition. A lower IC₅₀ value corresponds to a higher antioxidant capacity of plant extract (Patro *et al.*, 2005). The ability to scavenge the DPPH radical was calculated by:

$$I\% = \frac{A_0 - A_1}{A_0} \times 100$$

Where A₀ is the absorbance of the control at 30 min, and A₁ is the absorbance of the sample at 30 min.

Statistical analysis

All data were analyzed by variance analysis using SPSS 16. Experiments were tested using completely randomized design in factorial form (in three replicates). Mean comparisons were conducted using one way and univariate tests.

Results

Study of root morphology of inoculated plants indicated pink nodules on the roots of these plants.

A) Results of bacterial inoculation and SO₂ pollution on total antioxidants

Effect of bacterial inoculation

In this study, the total antioxidant activities (I %) and capacities (IC₅₀) of the samples were determined by DPPH-radical scavenging activity test. The results showed that bacterial inoculation had no significant effect on total antioxidant activity and capacity. Rhizobium inoculation did not cause stress conditions for plant, so the plant is normally and changes in the antioxidant system is not created.

Effect of SO₂ pollution

The DPPH radical scavenging activity and capacity indicated no significant change in 0 and 0.5 ppm concentrations of SO₂. Increasing SO₂ stress changed I% and IC₅₀ significantly. I% increased and IC₅₀ decreased in high concentration of SO₂ (1, 1.5 and 2 ppm) as compared with control plant. I% was at its highest value in 2 ppm of SO₂ with 80.58% increase in comparison with the controls (Fig. 1a). IC₅₀ was at its lowest value in 2 ppm with 44.13% decrease in comparison with the controls (Fig 1b).

Interaction effect of bacterial inoculation and SO₂ pollution

To understand the protective function of antioxidants against SO₂ stress, alfalfa plants were treated with native and standard *Rhizobium meliloti* followed by measurement of the level of I% and IC₅₀ (Table 1). Interaction effect of bacterial inoculation and SO₂ pollution on I% and IC₅₀ was significant as compared with control plants (no bacteri, no SO₂). Increasing SO₂ stress significantly increased I% and decreased IC₅₀ of alfalfa leaves compared to the control in the

experiment. Inoculation of alfalfa plant with native and standard *Rhizobium meliloti* significantly reduced the stress effects of high concentrations of SO₂ on I% and IC₅₀. Among bacterial inoculation and SO₂ pollution treatments, noninoculated plants under 2 ppm SO₂ (-R 2) and inoculated plants with native *Rhizobium meliloti*

under 0 ppm SO₂ (R_n 0) showed higher and lower level of I% respectively. I% was lower in inoculated treatments compared with control treatments. Interaction between bacterial inoculation and SO₂ pollution showed that they had significant effect ($p \leq 0.01$) on IC₅₀. Increasing SO₂ doses significantly decreased IC₅₀




Fig 1. (a) Antioxidant capacities (IC₅₀) and (b) antioxidant activities (I %) in the leaves of alfalfa under SO₂ pollution in 45 days plants. Data are mean \pm SE. Different letters indicate that the mean value is significantly different ($p < 0.01$).

The IC₅₀ in the leaves of noninoculated alfalfa plant under 2 ppm SO₂ (-R 2) was 54.24% lower than control treatments (-R0). The level of decreasing in IC₅₀ was slightly higher in inoculated plants. Interaction effect of bacterial inoculation and SO₂ pollution had shown that decline in IC₅₀ on noninoculated plants under 2 ppm SO₂ (-R2) was 54.24% but on inoculated plants with native and standard *Rhizobium meliloti* under 2 ppm SO₂ (R_n2 and R_s 2) was 45.93 and 46.16% respectively. IC₅₀ levels were lower in inoculated plants than noninoculated plants (Table 1).

B) Bacterial inoculation and SO₂ pollution on antioxidant activity (SOD, CAT, GPOX).

Effect of bacterial inoculation

The results showed that inoculation did not impose any significant effect on antioxidant activity (SOD, CAT, GPOX) and there was no significant difference among inoculated and noninoculated plants.

Effect of SO₂ pollution

There was significant difference among treatments under different concentrations SO₂ pollution in SOD, CAT and GPOX activity. The change in SO₂ concentration caused increased in antioxidant activity (Fig 2).

Superoxide dismutase activity

SOD activity was not different under the influence of 0 and 0.5 ppm of SO₂, but increased significantly with higher concentrations of SO₂. The SOD activity in the leaves of alfalfa at 1, 1.5 and 2 ppm of SO₂ pollution was 48.96%, 86.15% and 135.12% higher than 0 ppm of SO₂ pollution treatments, respectively (Fig. 2a).

Catalase activity

SO₂ pollution effect on CAT activity was significant. Compared with the control, CAT activity was affected by SO₂ treatments, was highest in the 2 ppm dose whereas being no change in other doses. The CAT activity in the leaves of alfalfa plant under 2 ppm 59.37% higher than control treatments (0 ppm) (Fig. 2b).

Guaiacol peroxidase activity

The antioxidant activity of GPOX was not different under the influence of 0 and 0.5 ppm of SO₂ pollution but increased in higher doses of SO₂ pollution. Among SO₂ pollution treatments, 2 ppm showed higher level of GPOX activity. The GPOX activity in the leaves of alfalfa at 1, 1.5 and 2 ppm of SO₂ pollution was 23.24, 57.73 and 77.91% higher than 0 and 0.5 ppm of SO₂ pollution treatments, respectively (Fig. 2c).




Fig 2. SOD (a), CAT (b) and GPOX(c) activity in the leaves of alfalfa under SO₂ pollution in 45 days plants. Data are mean ± SE. Different letters indicate that the mean value is significantly different ($p < 0.01$).

Table 1. Means comparison of interaction effect of bacterial inoculation (no-inoculation (-R), inoculation with native (R_n) and standard (R_s) *Rhizobium meliloti*) and SO₂ pollution (0, 0.5, 1, 1.5 and 2 ppm) on values of antioxidant capacities (IC₅₀) and antioxidant activities (I %) in 45 days plants. Similar words indicate not significantly different according to Duncan's test. The data are the means of three replicates ±SE and comparisons were performed separately for each index.

Bacterial inoculation	SO ₂ treatments (ppm)	IC ₅₀ (mg/ml)	I%
Noninoculation (-R)	0	17.44 ^a ± 0.27	7.21 ^g ± 0.11
	0.5	17.32 ^a ± 0.22	7.18 ^g ± 0.09
	1	13.33 ^c ± 0.20	9.38 ^e ± 0.14
	1.5	10.64 ^e ± 0.03	11.74 ^c ± 0.03
	2	7.98 ^g ± 0.04	15.66 ^a ± 0.13
Inoculation with native <i>Rhizobium</i> (R _n)	0	17.11 ^a ± 0.27	7.31 ^g ± 0.11
	0.5	16.99 ^a ± 0.12	7.35 ^g ± 0.05
	1	15.03 ^b ± 0.10	8.32 ^f ± 0.05
	1.5	11.63 ^d ± 0.04	10.74 ^d ± 0.04
	2	9.43 ^f ± 0.06	13.25 ^b ± 0.08
Inoculation with standard <i>Rhizobium</i> (R _s)	0	17.24 ^a ± 0.18	7.25 ^g ± 0.07
	0.5	17.30 ^a ± 0.16	7.23 ^g ± 0.06
	1	15.16 ^b ± 0.11	8.24 ^f ± 0.05
	1.5	11.64 ^d ± 0.10	10.74 ^d ± 0.09
	2	9.39 ^f ± 0.06	13.32 ^b ± 0.09

Interaction effect of bacterial inoculation and SO₂ pollution on antioxidant activity

Interaction of bacterial inoculation and SO₂ gas on antioxidant activity of SOD, CAT and GPOX was statistically significant. Combination of SO₂ pollution and bacterial inoculation had shown that the highest levels of SOD, CAT and GPOX activity were obtained at noninoculated plants under 2 ppm SO₂ (-R 2). Inoculated plants under high concentrations of SO₂ (1, 1.5 and 2 ppm)

showed antioxidant activity lower than noninoculated plant. For instance, noninoculated plant under 2 ppm SO₂ gas treatment showed 84.37% increase in the CAT activity but inoculated plant under 2 ppm SO₂ gas treatment showed 43.75% increase in the CAT activity. The rate of increase in antioxidant activity under SO₂ pollution stress in inoculated plant was lower than in the noninoculated plant (Fig 3).




Fig 3. Interaction effect of bacterial inoculation and SO₂ pollution on antioxidant activity of SOD (a), CAT (b) and GPOX (c) the leaves of alfalfa under SO₂ pollution in 45 days plants. Data are mean ± SE. Different letters indicate that the mean value is significantly different (*p* < 0.01).

Discussion

Various abiotic stresses lead to the overproduction of reactive oxygen species (ROS) in plants which are highly reactive and toxic and ultimately results in oxidative stress. However, the cells are equipped with excellent antioxidant defense mechanisms to detoxify the harmful effects of ROS. ROS are now also considered as key regulatory molecules vital for cells, but they cause cellular damage when produced in excess or when the antioxidant defense system is not properly functioning. Therefore, the concentration of ROS in cell must be controlled (Gill & Tuteja, 2010). After entering SO₂ to the leaf, oxidation of sulphite to sulphate occurs in the chloroplast. This oxidation gives rise to formation ROS such as O₂⁻ (Arora *et al.*, 2002; Li & Yi, 2012). When SO₂ is converted to sulphite in aqueous solution, it is a nucleophilic agent that is able to attack in numerous substrates by opening S—S bridges. This reaction, so called sulphitolysis, causes the inactivation of proteins like thioredoxins and, as a consequence, causes a severe reduction in plant growth or even cell death. Plants manage the internal sulphite concentration, i.e. they control the: (1) uptake of the gas by the laminar boundary layer, the cuticle or the guard cells and (2) rate of its metabolic conversion by feeding into sulphur assimilation stream for production of cysteine, or reoxidation into sulphate. Apoplastic peroxidases were described to detoxify sulphite. Non-enzymatic oxidation of sulphite, initiated by superoxide anions formed on the reduction site of the electron transport system in chloroplasts (Lang *et al.*, 2007). sulphite oxidation proceeds via a radical chain reaction involving light-dependent photosynthetic electron transport, sulphite oxidizing activity associated with isolated thylakoid membranes by the enzyme sulphite oxidase (SO) that the plant enzyme is localized in peroxisomes. Molecular oxygen as the terminal electron acceptor for plant SO and showed that it converts molecular oxygen into hydrogen peroxide (Hänsch *et al.*, 2006). SO plays a key role in protection of plants from the damaging effects of SO₂. SO is a housekeeping enzyme that oxidize sulphite using H₂O₂ and different phenolic compounds. It is involved in the recycling of sulphur. It may also protect the thioredoxin system from damage (Lang *et al.*, 2007). In such conditions, plants develop a high efficient antioxidant enzymatic defense system to

increase tolerance to different stress factors (Gill & Tuteja, 2010).

In this study, values of IC₅₀, I% and antioxidant activity indicated no significant difference at 0.5 ppm of SO₂ as compared with control plants (exposed to 0 ppm) because may be stress conditions are not created in 0.5 ppm of SO₂, but indicated significant differences in higher concentration. In higher concentrations of SO₂ (1, 1.5 and 2 ppm), IC₅₀ value decreased with increasing stress intensity but I% increased. Increasing of I% means more antioxidants has been produced with increasing stress intensity. Increase of DPPH-radical scavenging activity has been reported in many studies. DPPH-radical scavenging activity significantly increased as compared with control plants in *Oryza sativa* seedlings under heat shock treatment (Kang & Saltveit, 2002) and in *Cakile maritima* exposed to salinity stress (Ksouri *et al.*, 2007). Use of silicon and salinity on *Sorghum bicolor* increased DPPH-radical scavenging activity as compared with control plants (Kafi *et al.*, 2011).

Guaiacol peroxidase (GPOX) decomposes indole-3-acetic acid (IAA) and has a role in the biosynthesis of lignin and defence against biotic stresses by consuming H₂O₂ (Gill & Tuteja, 2010). The activity of GPOX varies considerably depending upon plant species and stresses condition. In this study, GPOX activity increased in 1, 1.5 and 2 ppm of SO₂ as observed in *Zizyphus mauritiana* and *Mangifera indica* (Rao & Dubey, 1990) exposed to SO₂ pollution.

Catalases (CAT) are tetrameric heme containing enzymes with the potential to directly dismutate H₂O₂ into H₂O and O₂ and is indispensable for ROS detoxification during stressed conditions. CAT has one of the highest turnover rates for all enzymes: one molecule of CAT can convert 6 million molecules of H₂O₂ to H₂O and O₂ per minute. CAT is important in the removal of H₂O₂ generated in peroxisomes by oxidases involved in β-oxidation of fatty acids, photorespiration and purine catabolism (Gill & Tuteja, 2010). In this study, CAT activity indicated no significant difference at 0.5, 1 and 1.5 ppm concentration of SO₂ as compared with control plants but in 2 ppm of SO₂ indicated a significant increase. CAT is an antioxidant that is activated in severe stress conditions (Gill & Tuteja, 2010). Various studies have reported different results from the CAT activity. Increase in CAT activity indicated in

wheat plant under drought stress (Simova- Stoilova *et al.*, 2010). Glutathione S-transferase (GST) is a cytosolic enzyme that is induced by various stressors and is involved in the detoxification of xenobiotics and in the conjugation of drugs and carcinogens. It is also involved in the detoxification of reactive oxygen species (ROS) and in the regulation of cell growth and differentiation. The activity of GST is increased in plants under stress conditions, and it is considered to be a marker of plant stress tolerance. In this study, the activity of GST was measured in faba bean plants under SO₂ pollution. The results showed that the activity of GST increased significantly in plants treated with 1, 1.5 and 2 ppm of SO₂ compared to control plants. This increase in GST activity is likely due to the increased production of ROS under stress conditions, which leads to the activation of GST. The increase in GST activity is also consistent with the increased activity of other antioxidant enzymes, such as SOD and GPOX, observed in this study. These results suggest that the increased activity of GST is a response to the oxidative stress induced by SO₂ pollution. The increased activity of GST may help to detoxify ROS and other harmful substances, thereby reducing the damage to plant cells and improving plant growth and yield under stress conditions. The results of this study also suggest that the inoculation of faba bean plants with native and standard *R. meliloti* can improve their tolerance to SO₂ pollution. This is likely due to the increased activity of antioxidant enzymes, such as SOD, GPOX, and GST, induced by the inoculation. The increased activity of these enzymes may help to reduce the damage to plant cells and improve plant growth and yield under stress conditions. The results of this study also suggest that the inoculation of faba bean plants with native and standard *R. meliloti* can improve their tolerance to SO₂ pollution. This is likely due to the increased activity of antioxidant enzymes, such as SOD, GPOX, and GST, induced by the inoculation. The increased activity of these enzymes may help to reduce the damage to plant cells and improve plant growth and yield under stress conditions.

In this study, inoculation with native and standard *R. meliloti* had no significant effects on values of IC₅₀, I%, GPOX, SOD and CAT activity, as reported by Gabballah & Gomaa (2005). Their study showed that inoculation of two cultivars of *Vicia faba* with *Rhizobium* had no significant effects on SOD activity.

Interaction of inoculation and SO₂ treatment in this study was significant. Indeed *Rhizobium* inoculation under SO₂ condition showed significant effect on the values of IC₅₀, I%, GPOX, SOD and CAT activity. Rhizobacteria such as *Rhizobium* can promote plant growth and by mechanisms that used for this growth promotion can reduce stress conditions for plants. Plant growth promotion by rhizobacteria can occur directly and indirectly. There are several ways by which PGPR can affect plant growth directly, e.g. by fixation of atmospheric nitrogen, solubilization of minerals such as phosphorus, production of siderophores that solubilize and sequester iron, or production of plant growth regulators (hormones) that enhance plant growth at various stages of development. Indirect growth promotion occurs when PGPR promote plant growth by improving growth restricting conditions. This can happen directly by producing

resistance to biotic and abiotic stresses (Timmusk, 2003). Indeed can be concluded that bacterium has been reduced antioxidant activity and capacity by reducing the effects of high concentrations of SO₂ and stress conditions. Different studies have expressed different conclusions about the interaction between bacterial inoculation and stress. Inoculation of Lettuce under salinity stress with *Rhizobium* sp. and *Serratia* sp. decreased enzyme activity, including glutathione reductase and ascorbate peroxidase, with increasing salinity stress (Han & Lee, 2005). A clear decline in SOD activity in two cultivars of faba bean was observed with increasing salinity stress. Use of *Rhizobium* inoculation and sodium benzoate increased SOD activity in faba bean plants under salinity (Gabballah & Gomaa, 2005).

Conclusion

The induction of the cellular antioxidant machinery is important for protection against ROS. Overexpression of ROS scavenging enzymes like SOD, CAT, and GPOX resulted in abiotic stress tolerance in various crop plants due to efficient ROS scavenging capacity. Low concentration of SO₂ (0.5 ppm) doesn't create stress conditions in alfalfa therefore activity and capacity of most of antioxidants don't alter in this concentration. In higher concentrations of SO₂ (1, 1.5 and 2 ppm), activity and capacity of alfalfa antioxidants increase with increasing stress intensity. Inoculation of alfalfa plant with native and standard *Rhizobium meliloti* significantly reduced the negative effects of high concentration of SO₂ on antioxidants activity and capacity.

References

- Abe N, Murata T, Hirota A. 1998. Novel 1,1-diphenyl-2-picrylhydrazyl radical scavengers, bisorbicillin and demethyltrichodimerol, from a fungus. *Bioscience, Biotechnology and Biochemistry* 62: 61-662.
- Aeini M, Yousefi Rad M, Ehteshami SM. 2012. The effects of growth stimulant bacteria on qualitative and quantitative yield of *Aloe vera*. *Annals of Biological Research* 3 (12): 5669-5673.
- Agrawal M, Nandi, PK, Rao DN. 1985. Effects of sulphur dioxide fumigation on soil system and

- growth behaviour of *Vicia faba* plants. *Plant*
 Arora A, Sairam RK, Srivastava GC. 2002. Oxidative stress and antioxidative systems in plants. *Current Science* 82: 1227–1238
- Bai Y, Zhou X, Smith DL. 2003. Crop ecology, management and quality: enhanced soybean plant growth resulting from coinoculation of *Bacillus* strains with *Bradyrhizobium japonicum*. *Crop Science* 43: 1774-1781.
- Bashan Y, Levanony H, Mitiku G. 1989. Changes in proton efflux of intact wheat roots induced by *Azospirillum brasilense* Cd. *Canadian Journal of Microbiology* 35: 691-697.
- Benabderrahim MA, Mansour H, Ferchichi A. 2009. Diversity of Lucerne (*Medicago sativa* L.) Populations in South Tunisia. *Pakistan Journal of Botany* 41: 2851-2861.
- Bernardi R, Nali C, Gargiulo R, Pugliesi C, Lorenzini G, Durante M. 2001. Protein pattern and Fe-superoxide dismutase activity of bean plants under sulphur dioxide stress. *Journal of Phytopathology* 149: 477-480.
- Caetano-Anolles G, Wall LG, De Micheli AT, Macchi EM, Bauer WD, Favelukes G. 1988. Role of motility and chemotaxis in efficiency of nodulation by *Rhizobium meliloti*. *Plant Physiology* 86: 1228-1235.
- Cakmak I, Marschner H. 1992. Manganese deficiency and high light intensity enhance activities of superoxide dismutase, ascorbate peroxidase, and glutathione reductase in bean leaves. *Plant Physiology* 98: 1222-1227.
- Chaparzadeh N, Amico MLD, Khavari-Najad RA, Navarizzo F. 2004. Antioxidative responses of *Calendula officinalis* under salinity conditions. *Plant Physiology and Biochemistry* 42: 695-701.
- Dimkpa C, Weinand T, Asch F. 2009. Plant-rhizobacteria interactions alleviate abiotic stress conditions. *Plant, Cell and Environment* 32:1682-1694.
- Estefan M, Munteanu, N, Stoleru, V, Mihasan, M. 2013. Effects of inoculation with plant growth promoting rhizobacteria on photosynthesis, antioxidant status and yield of runner bean. *Romanian Biotechnological Letters* 18: 8132-8143.
- Gabballah MS, Gomaa AM. 2005. Interactive effect of *Rhizobium* inoculation, sodium benzoate and salinity on performance and oxidative stress in two fababean varieties. *International Journal of Agriculture and Biology* 7(3): 495-498.
- and Soil 86(1): 69-78.
- Gentili F, Jumpponen A. 2006. Potential and possible uses of bacterial and fungal biofertilizers (chapter 1). In: Rai MK, ed.: *Handbook of microbial biofertilizers*. New York: Haworth press, 1-28.
- Giannopolitis CN, Ries SK. 1977. Superoxide Dismutases: I. occurrence in higher plants. *Plant Physiology* 59: 309-314.
- Gill SS, Tuteja N. 2010. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiology and Biochemistry* 48: 909-930.
- Gostin IN. 2009. Air pollution effects on the leaf structure of some Fabaceae species. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* 37 (2): 57-63.
- Han HS, Lee KD. 2005. Plant growth promoting rhizobacteria effect on antioxidant status, photosynthesis, mineral uptake and growth of *Lettuce* under soil salinity. *Research Journal of Agriculture and Biological Sciences* 1(3): 210-215.
- Hänsch R, Lang C, Riebeseel E, Lindigkeit R, Gessler A, Rennenberg H, Mendel R.R. 2006. Plant sulfite oxidase as novel producer of H₂O₂: Combination of enzyme catalysis with a subsequent non-enzymatic reaction step. *Journal of Biological Chemistry* 281:6884-6888.
- Kafi M, Nabati J, Masoumi A, Mehrgerdi MZ. 2011. Effect of salinity and silicon application on oxidative damage of sorghum [*Sorghum bicolor* (L.) monech.]. *Pakistan journal of Botany* 43(5): 2457-2462.
- Kang HM, Saltveit ME. 2002. Antioxidant enzymes and DPPH-radical scavenging activity in chilled and heat-shocked rice (*Oryza sativa* L.) seedlings radicles. *Agricultural and Food Chemistry* 50: 513-518.
- Ksouri R, Megdiche W, Debez A, Falleh H, Grignon C, Abdelly C. 2007. Salinity effects on polyphenol content and antioxidant activities in leaves of the halophyte *Cakile maritima*. *Plant Physiology and Biochemistry* 45: 244-249.
- Lang C, Popko J, Wirtz M, Hell R, Herschbach C, Kreuzwieser J, Rennenberg H, Mendel R, Hansch R. 2007. Sulphite oxidase as key enzyme for protecting plants against sulphur dioxide. *Plant, Cell and Environment* 30:447-455.
- Li L, Yi H. 2012. Effect of sulfur dioxide on ROS production, gene expression and antioxidant

- enzyme activity in Arabidopsis plants. *Plant Physiology and Biochemistry* 58: 46-53.
- Martínez-Viveros O, Jorquera MA, Crowley DE, Gajardo G, Mora ML. 2010. Mechanisms and practical considerations involved in plant growth promotion by rhizobacteria. *Journal of Soil Science Plant Nutrition* 10(3): 293-319.
- Millner PD, Kitt DG. 1992. The Beltsville method of soilless production of vesicular-arbuscular mycorrhizal fungi. *Mycorrhiza* 2: 9-15.
- Moini L, Fani A, Bakhtyar M, Rafeie M. 2011. Correlation between the concentration of air pollutants (CO, SO₂ and NO₂) and pulmonary function. *Journal of Shahrekord University of Medical Sciences* 13(1): 27-35.
- Molla AH, Shamsuddin ZH, Halimi MS, Morziah M, Puteh AB. 2001. Potential for enhancement of root growth and nodulation of soybean co-inoculated with *Azospirillum* and *Bradyrhizobium* in laboratory systems. *Soil Biology and Biochemistry* 33: 457-463.
- Patro BS, Bauri AK, Mishra S, Chattopadhyay S. 2005. Antioxidant activity of *myristica malabarica* extracts and their constituents. *Journal of Agricultural and Food Chemistry* 53(17): 6912-6918.
- Polle A, Otter T, Seifert F. 1994. Apoplastic peroxidases and lignification in needles of norway spruce (*Picea abies* L.). *Plant Physiology* 106: 53-56.
- Rai R, Rajput M, Agrawal M, Agrawal SB. 2011. Gaseous air pollutants: a review on current and future trends of emissions and impact on agriculture. *Journal of Scientific Research Banaras Hindu University, Varanasi* 55: 77-102.
- Rao MV, Dubey PS. 1990. Explanations for the differential response of certain tropical tree species to SO₂ under field conditions. *Water, Air and Soil Pollution* 51: 297-305.
- Sadovinkova YN, Bepalova LA, Antonyuk LP. 2003. Wheat gram agglutinin is a growth factor for bacterium *Azospirillum brasilense*. *Doklady Biochemistry and Biophysics* 398:103-105.
- Salehi M, Salehi F, Poustini K, Heidari-Sharifabad H. 2008. The effect of salinity on the nitrogen fixation in 4 cultivars of *Medicago sativa* L. in the seedling emergence stage. *Research Journal of Agriculture and Biological Sciences* 4(5): 413-415.
- Sha C, Wang T, Lu J. 2010. Relative sensitivity of Wetland plants to SO₂ pollution. *Wetlands* 30: 1023- 1030.
- Saharan BS, Nehra V. 2011. Plant growth promoting rhizobacteria: A critical review. *Life sciences and medicine research*. Available at <http://astonjournals.com/lsmr-21>
- Simova-Stoilova L, Vaseva I, Grigorova B, Demirevska K, Feller U. 2010. Proteolytic activity and cysteine protease expression in wheat leaves under severe soil drought and recovery. *Plant Physiology and Biochemistry* 48: 200-206.
- Smith SJ, Aardenne JV, Klimont Z, Andres RJ, Volke A, Delgado Arias S. 2011. Anthropogenic sulfur dioxide emissions: 1850–2005. *Atmospheric Chemistry and Physics* 11: 1101–1116.
- Swanepoel JW, Kruger GHJ, Heerden PDR. 2007. Effects of sulphur dioxide on photosynthesis in the succulent *Augeacapensis* Thunb. *Journal of Arid Environments* 70: 208-221.
- Swift M, Bignell D. 2001. Standard methods for assessment of soil biodiversity and land use practice. Available at: <http://www.icraf.cgiar.org/sea>.
- Tank N, Saraf M. 2010. Salinity-resistant plant growth promoting rhizobacteria ameliorates sodium chloride stress on tomato plants. *Journal of Plant Interactions* 5: 51-58.
- Timmusk S. 2003. Mechanism of action of the plant growth promoting bacterium *Paenibacillus polymyxa*. *Acta Universitatis Upsaliensis, Uppsala, Sweden*.
- Wali B, Iqbal M, Mahmooduzzafar. 2007. Anatomical and functional responses of *Calendula officinalis* L. to SO₂ stress as observed at different stages of plant development. *Flora* 202: 268–280.
- Wang YX, Oyaizu H. 2009. Evaluation of the phytoremediation potential of four plant species for dibenzofuran-contaminated soil. *Journal of Hazardous Materials* 168: 760-764.

