

Chromosome C-banding in *Mus musculus* L.1766 strains shows a fixed position for the centromere and variable amounts in different populations

Farahnaz Molavi¹, Jamshid Darvish^{*1,2}, Farhang Haddad¹ and Maryam M. Matin^{1,3}

¹Department of Biology, Faculty of Science, Ferdowsi University of Mashhad, Mashhad, Iran

²Rodentology Research Department, Ferdowsi University of Mashhad, Mashhad, Iran

³Cell and Molecular Biotechnology Research Group, Institute of Biotechnology, Ferdowsi University of Mashhad, Mashhad, Iran

*Corresponding author: darvishj2001@yahoo.com

Received: 13/07/2015 Accepted: 25/10/2015

Abstract

Three subspecies of *Mus musculus* have been recognized in Iran so far. The house mouse (genus *Mus*, species *Mus musculus*) are recognized for their highly conserved morphology and chromosomal structure, but some chromosomal characters offer accurate taxonomic markers in this species that has been shown any unambiguous diagnostic morphological traits. Among the chromosomal characters, centromeric heterochromatin is more useful to identify mouse subspecies and populations. In this study, Samples were collected from 27 stations in Iran and study was performed by the centromeric heterochromatin banding (C-banding). Results indicated that all samples had 40 acrocentric chromosomes and all chromosomes had fixed position for the centromere. The strains had the same amount of C-banding material on homologous chromosomes but showed variation in the amount on different populations.

Key words: Rodentia; Muridae; Karyology; House mouse; Karyotype; C- Banding; Iran

نواربندی سانترومیری گستره های موش های خانگی (*Mus musculus* L.1766) ایران، نشان دهنده موقعیت ثابت سانترومر با مقادیر متغییر هتروکروماتین سانترومیری در جمعیت های مختلف است

فرحناز مولوی^۱، جمشید درویش^{۱،۲*}، فرهنگ حداد^۱، مریم مقدم متین^{۳،۱}

^۱دانشگاه فردوسی مشهد، دانشکده علوم، گروه زیست شناسی

^۲دانشگاه فردوسی مشهد، گروه پژوهشی جوندگان شناسی

^۳دانشگاه فردوسی مشهد، گروه پژوهشی بیوتکنولوژی مولکولی و سلولی

چکیده

مدت ها است که سه زیرگونه از گونه موش خانگی در ایران شناسایی شده اند. موش خانگی (جنس *Mus*، گونه *M. musculus*) با ساختار کروموزومی و ریختی پایدار خود شناخته می شوند. اما برخی از صفات کروموزومی به عنوان نشانگرهای مورد اعتماد در این گونه شناخته شده اند. در میان این صفات کروموزومی، ساختار هتروکروماتین سانترومیری جز این نشانگرها است که برای شناسایی جمعیت ها و زیرگونه های موش خانگی استفاده شده است. در این مطالعه، ۲۷ جمعیت از ایران مورد بررسی قرار گرفتند و ساختار هتروکروماتین آنها با هم مقایسه شد. نتایج نشان داد که در تمام افراد مورد تحقیق سانترومر جایگاه یکسانی در کروموزوم ها دارد. و با اینکه اندازه هتروکروماتین سانترومیری در بین کروموزوم های همولوگ یکسان است ولی بین جمعیت های مختلف تفاوت اندازه دارند.

واژه های کلیدی: جوندگان؛ موش؛ کاریولوژی؛ موش خانگی؛ کاریوتایپ؛ ایران

Introduction

Muridae species despite high morphologic similarity display much higher variability of karyotypes (Romanenko *et al.*, 2007). The centromeric domain of most mammalian chromosomes contains repeated DNA sequences (Britten and Kohne, 1968). In the mouse, *Mus musculus*, the major repetitive DNA component is the satellite DNA sequence family first isolated by Kit (1961). As shown by Pardue and Gall (1969) and Jones (1970) this major satellite is present on all *M. musculus* chromosomes with the exception of the Y chromosome, and constitutes around 10% of the *M. musculus* genome. In 1983 Pietras *et al.* isolated a second repetitive DNA family from *M. musculus*, which comprised a much smaller proportion (<1%) of the genome and called it the minor satellite. By *in situ* hybridization Wong and Rattner (1988), Joseph *et al.* (1989) and Broccoli *et al.* (1990) have shown that the minor satellite DNA sequences within *M. musculus* appear to be physically very close to the primary constriction (*i.e.* the centromere) in these chromosomes. In comparison, the major satellite DNA sequences, although localized to the pre-centromeric heterochromatin in *M. musculus*, occupy a separate non-overlapping domain to that of the minor satellite DNA sequences (Joseph *et al.*, 1989). This domain, together with its associated C-band, can be deleted from a mouse chromosome without affecting centromere function, which argues strongly that major satellite DNA sequences are not directly involved in centromere function (Broccoli *et al.*, 1990).

Materials and Methods

40 specimens were captured from 27 stations of Iran (Table 1 and Fig. 1). Sample codes of specimens are available in the Rodentology Research Department of Ferdowsi University of Mashhad, Iran. Chromosome spreads were obtained from bone marrow cells according to Yosida (1973). About 50 to 100 metaphase plates from both male and female specimens were examined and at least 30 good chromosomal spreads were photographed using a 100x zoom digital CCD camera. The karyological characteristics of all specimens were prepared by Karyological Analysis software (version 1.2, 2010). Chromosomes were classified according to Levan (Levan *et al.*, 1964), and each chromosome

was placed next to its presumed homologue to determine the diploid chromosome number (2n). C-banding: Barium hydroxide - Saline - Giemsa (BSG) method was performed according to Summer (1972).




Figure 1. Map of specimen collection sites for *M. musculus* strains in Iran (black dots are stations)

Results

In this study, 40 specimens of *M. musculus* including all 3 subspecies present in Iran were studied. The karyological characteristics of these species are described in Figure 2. In most of the strains every chromosome except the Y chromosome was found to have C-banding material (Fig. 3). The size of the C-banding region was the same for both homologous chromosomes in each strain (Figure 4), but the size of this region on some chromosomes varied from one population to another (Fig. 5 and 6). In Mashhad populations, the amount was approximately the same on each chromosome and we have used this strain as a standard for comparison. The centromere region size was variable in populations. For example, in Mashhad population, chromosomes 8 and 14 had smaller amounts of C-banding material than Gonabad population; whereas, chromosomes 3 and 19 had larger amounts of that (Fig. 4). Moreover, the Zahedan strain differs from Zabol in having very little C-banding material on chromosome 10 and 13 and

having larger amount of C-banding material on chromosomes 5 and 12.

Table 1. Sampling localities, geographical coordinates, and sample codes of *M. musculus* from different localities of Iran.

Row	Taxon	Locality	Longitude	Latitude	2n
1	<i>M. m. bacterianus</i>	Kerman	30° 17' 46" N	57° 05' 23" E	40
2	<i>M. m. musculus</i>	Gonabad	34° 21' 10" N	58° 41' 1" E	40
3	<i>M. m. musculus</i>	Torbate Jam	35° 14' 38" N	60° 37' 21" E	40
4	<i>M. m. musculus</i>	Sarakhs	36° 32' 42" N	61° 9' 28" E	40
5	<i>M. m. musculus</i>	Dargaz	37° 26' 40" N	59° 6' 29" E	40
6	<i>M. m. musculus</i>	Kalat	36° 59' 33.01" N	56° 45' 23.83"E	40
7	<i>M. m. Isatissus</i>	Isfahan	32° 63' 35" N	51° 65' 36" E	40
8	<i>M. m. bacterianus</i>	Zahedan	29° 29' 47" N	60° 51' 46" E	40
9	<i>M. m. bacterianus</i>	Khash	28° 13' 16" N	61° 12' 57" E	40
10	<i>M. m. bacterianus</i>	Zabol	31° 1' 43" N	61° 49' 4" E	40
11	<i>M. m. domesticus</i>	Chabahar	25° 17' 31" N	60° 64' 35" E	40
12	<i>M. m. Isatissus</i>	Shiraz	29° 61' 0" N	52° 54' 0" E	40
13	<i>M. m. Isatissus</i>	Yazd	31° 53' 50" N	54° 22' 4" E	40
14	<i>M. m. musculus</i>	Esfarayen	37° 73' 03" N	57° 50' 72" E	40
15	<i>M.m. domesticuss</i>	Eizeh	31° 50' 48" N	49° 50' 36" E	40
16	<i>M. m. domesticuss</i>	Mamasani	30° 7' 0" N	51° 31' 0" E	40
17	<i>M. m. bacterianus</i>	Sabzevar	36° 70' 35" N	59° 6' 44" E	40
18	<i>M. m. musculus</i>	Tabadkan	36° 29' 0" N	59° 40' 58" E	40
19	<i>M. m. musculus</i>	Mashhad	36° 18' 0" N	59° 36' 0" E	40
20	<i>M. m. bacterianus</i>	Rask	26° 13' 47" N	61° 13' 4" E	40
21	<i>M. m. bacterianus</i>	Birjand	32° 87' 0" N	59° 20' 0" E	40
22	<i>M. m. bacterianus</i>	Qasregand	26° 15' 60" N	60° 45' 14" E	40
23	<i>M. m. bacterianus</i>	Negor	25° 38' 15" N	61° 12' 33" E	40
24	<i>M. m. musculus</i>	Kardeh	36° 16' 51" N	59° 34' 59" E	40
25	<i>M. m. bacterianus</i>	Zahedan2	37° 33' 19" N	45° 04' 21" E	40
26	<i>M. m. musculus</i>	Neyshabour	36° 20' 48" N	58° 10' 48" E	40
27	<i>M. m. domesticus</i>	Bandar Mahshahr	30° 54' 32" N	49° 11' 58" E	40

longest/shortest=3.1756
Number of chromosome which (long arm/short arm)>2: 40 (1.0000)
The karyotype asymmetry index (Arano, 1963), As K%=100.00%
The total form percent (Huziwara, 1962), TF%=0.00%
The index of karyotype symmetry (Greilhuber and Speta, 1976), Syi=0.00%
The index of chromosomal size resemblance (Greilhuber and Speta, 1976), Rec=61.69%
The intra chromosomal asymmetry index (Romero Zarco, 1986), A1 = 1.00
The inter chromosomal asymmetry index(Romero Zarco, 1986), A2=0.26
The degree of asymmetry of karyotype (Watanabe et al., 1999), A=1.00
The dispersion index (Lavanaia and Srivastava, 1992), DI=0.00
The asymmetry index (Paszko, 2006), AI=0.00
Cytotype:4B

Figure 2. A example of the Karyological Analysis software output of *Mus musculus*.




Figure 3. Karyotype of *M. musculus* in Mashhad (a), Yazd (b), Zahedan (c) and Bandar Mahshahr (d)



Figure 4. The same size of the C-banding region of *Mus musculus* for both homologous chromosomes in Mashhad strain.




Figure 5. The size of the C-banding regions on chromosomes of *M. musculus* strains captured in Mashhad (left) and Gonabad (right).




Figure 6. The size of the C-banding region of *Mus musculus* for Zahedan (left) and Zabol (right) chromosomes .

Discussion

Chromosomal evolution in house mouse show reticulated evolution and speciation (Gunduz, 2010). Sub species of *M. musculus* are parapatric (Boursot *et al.*, 1989) Despite the reduction of meiotic fitness in hybrids (Sharma *et al.*, 2003), We Have been reported many hybrids of them in contact zones (Guenet, 2003). Information about highly polymorphic minisatellite markers suggested past and present genetic exchanges among house mouse subspecies (Bonhomme *et al.*, 2007). Rajabi-mahan, *et al.* (2012) showed *Mus musculus castaneus* is a polytypic subspecies of this species. In 2006, a transition zone of the house mouse have been reported from eastern Iranian plateau by Darvish in 2006. Malcon *et al.* in 2007 showed chromosomal characters are very useful for recognizing species in rodentia. The primary and secondary constrictions have already been used for comparison between populations in mice by Dev *et al.* (1971). They showed that almost all chromosomes could be distinguished by the presence or absence of large secondary constrictions located near to the C-banding region. In 1971 Eicher used the variant amount of secondary constriction region on chromosome 19 for calculating the degree of diversity in *Mus musculus*. He found that almost every chromosome of *M. m. mollossinus* was different from that of *M. musculus* in the size of C-banding region. Since *M. m. mollossinus* is interfertile with *M. musculus*, markers for every mouse chromosome except the Y chromosome are potentially available. The C-banding regions of mouse chromosomes are composed mainly of highly repetitious DNA, called satellite DNA (Pardu and Gall 1970). It is consisted of millions of copies of very short nucleotide sequences, or minor variants of them (Southerne, 1970). Polymorphic variants have been reported within the chromosomes of laboratory mice as well as in different species of wild mice (Forejt, 1973). Dev *et al.* (1973) looked at C-band variants within *M. musculus* strains and differences in the constrictions of chromosomes in these species using quinacrine fluorescence. Both Dev *et al.* (1973) and Davidson (1989) referred to a C-band variant on chromosome 1, although no reference was made to the type of DNA sequences involved. Vig and Richards (1992) have argued that the formation of the primary constriction in some mouse chromosomes (chromosome no. 1) does

not always require the presence of mouse minor satellite DNA sequences. Our results show that the amount of C-banding material on each chromosome is characteristic for each population of *Mus musculus*. The existence of obvious differences in the distribution of C-banding material among the chromosomes in different populations, suggests that the amount of C-banding material on a chromosome is a polymorphic trait that is inherited in a simple fashion.

Acknowledgements

We are grateful to Mortaza Radmanesh for his generous help and M. Mohammadyari, M. Habbiby, A. Khajea and A. Khosravi for their efforts in collecting the specimens. This study was supported by a grant (no. 27137) from Ferdowsi University of Mashhad.

References

- Bonhomme F, Rivals E, Orth A, Grant G, Jeffreys AJ, Rj Bois, P. 2007. Species-wide distribution of highly polymorphic minisatellite markers suggests past and present genetic exchanges among house mouse subspecies. *Genome Biol* 8: R80.
- Boursot P, Bonhomme F, Catalan J, Moriwaki K. 1989. Variations of Y chromosome repeated sequence across subspecies of *Mus musculus*. *Heredity* 63:289-297.
- Britten RJ, and Kohne DE. 1968. Repeated DNA sequences in DNA. *Science* 161: 529-540.
- Broccoli D, Miller OJ, Miller DA. 1990. Relationship of minor satellite to centromere activity. *Cytogenet Cell Genet* 54: 182-186.
- Darvish J, Orth A, Bonhomme F, 2006. Genetic transition in the house mouse *Mus musculus* of eastern Iranian plateau. *Folia Zool* 55: 176-182.
- Davidson MT. 1989. Centromeric chromosome variants. In *Genetic Variants and Strains of the Laboratory Mouse*, 2nd edi (ed. M. F. Lyon and A. G. Searle), Oxford University Press.
- Dev VG, Grewal MS, Miller DA, Kouri RE, Hutton JJ, Miller OJ. 1971. The quinacrine fluorescence karyotype of *Mus musculus* and demonstration of strain differences in secondary constrictions. *Cytogenet Genome Res* 10: 436-451.

- Dev VG, Miller DA, Miller OJ. 1973. Chromosome markers in *Mus musculus* strain differences in C-banding. *Genetics* 75: 663-670.
- Eicher EM. 1971. The identification of the chromosome bearing linkage group XII in the mouse. *Genetics* 69: 267-271.
- Flamm WG. 1972. Highly repetitious sequences of DNA in chromosomes. *Int Rev Cytol* 32:1-51.
- Forejt F. 1973. Centromeric heterochromatin polymorphism in the house mouse. *Chromosoma* 43: 187-201.
- Jones K. 1970. Chromosomal and nuclear location of mouse satellite DNA in individual cells. *Nature* 225: 912-915.
- Joseph A, Mitchell AR, Miller OJ. 1989. The organisation of the mouse satellite DNA at centromeres. *Exp Cell Res* 183: 494-500.
- Kit S, 1961. Equilibrium sedimentation in density gradients of DNA preparations from animal tissues. *J Mol Biol* 3: 711-716.
- Malcon A, Smith F, Trifonor V. 2007. Mammalian karyotype evolution. *Nature* 63: 950-961.
- Pardue ML, Gall JG. 1969. Molecular hybridisation of radioactive DNA to the DNA of cytological preparations. *Proc Nat Acad Sci USA* 64: 600-604.
- Pietras DF, Bennett KL, Siracusa LD, Woodworth-Gutai M, Chapman VM, Gross KW, Kane-Haas C, Hastie ND. 1983. Construction of a small *Mus musculus* repetitive DNA library: identification of a new satellite sequence in *Mus musculus*. *Nucleic Acids Res* 11: 6965-6983.
- Guenet JL, Bonhomme F. 2003. Wild mice an ever increasing contribution to a popular mammalian model. *Trends genet* 19: 5467-5472.
- Gunduz I. 2010. Staggered chromosomal hybrid zones in the house mouse: Relevance to reticulate evolution and speciation. *Genes* 1: 193-209.
- Rajabi-mahan H, Orth A, Siah sarvie R, Boursot P, Darvish J, Bonhomme F, 2012. The south-eastern house mouse *Mus musculus castaneus* (Rodentia: Muridae) is a polytypic subspecies. *Biol J Linn Soc Lond* 107: 295-306.
- Romanenko SA, Volobouev VT, Perelman PL, Lebedev VS, Serdukova NA, Trifonov VA, Biltueva LS, Nie V, Obrien PCM, Bulatova NS, Ferguson-Smith MA, Yang F, Graphodatsky AS. 2007. Karyotype evolution and phylogenetic relationships of hamsters (Cricetidae, Muroidea, Rodentia) inferred from chromosomal painting and banding comparison. *Chromosome Res* 15: 283-297.
- Sharma T, Bardhan A, Bardhan M. 2003. Reduced meiotic fitness in hybrids with heterozygosity for heterochromatin in the speciating *Mus Tricolor* complex. *Bioscience* 28: 137-143.
- Southerne M. 1970. Base sequence and evolution of guinea pig a-satellite DNA. *Nature* 227: 794-798.
- Vig BK, Richards BT. 1992. Formation of primary constriction and heterochromatin in mouse does not require minor satellite DNA. *Exp Cell Res* 201: 292-298.
- Wong AK, Rattner JB, 1988. Sequence organisation and cytological localization of the minor satellite of mouse. *Nucleic Acids Res* 16: 11645-11661.